arXiv:1212.5319v1 [hep-th] 21 Dec 2012

Preprint typeset in JHEP style - HYPER VERSION

Holographic Brownian Motion in Two Dimensional Rotating Fluid

Ardian Nata Atmaja[1,2]

[1]*Research Center for Physics, Indonesian Institute of Sciences (LIPI)*
Kompleks PUSPITEK Serpong, Tangerang 15310, Indonesia.
[2]*Indonesia Center for Theoretical and Mathematical Physics (ICTMP)*
Bandung 40132, Indonesia.
email: `ardian@teori.fisika.lipi.go.id`

Abstract: The Brownian motion of a heavy quark under a rotating plasma corresponds to BTZ black hole is studied using holographic method from string theory. The heavy quark represented as the end of string at the boundary of BTZ black hole and the corresponding rotating plasma is two dimensional spacetime. The string fluctuation requires the angular velocity to be equal to the ratio between inner horizon and outer horizon, known as terminal velocity and also related to the zero total force condition. With this angular velocity, the string fluctuation solution has oscillatory modes in time and radial coordinates. We show the displacement square of this solution behaves as a Brownian particle in non-relativistic limit. For relativistic case, we argue that it is more appropriate to compute just the the leading order of low frequency limit of random-random force correlator. The Brownian motion relates this correlator with physical observables: mass of Brownian particle, friction coefficient and temperature of the plasma.

Keywords: Brownian motion, BTZ black hole, rotating plasma.

Contents

1. Introduction **1**

2. Gravity Setup **3**

3. Small Fluctuation **4**
  3.1 Trivial Solution 4
  3.2 Linear Solution 5

4. Displacement Square **7**
  4.1 Hawking Radiation 7
  4.2 Position of the Brownian particle 9

5. Random-Random Force Correlator **11**
  5.1 Non-rotating BTZ black hole 12
  5.2 Rotating BTZ black hole 13

6. Discussion and Conclusion **14**

A. Non-relativistic Brownian Motion **16**

1. Introduction

AdS/CFT correspondence has been widely used to solve some problems in non-perturbative gauge theories. Some of the applications of AdS/CFT correspondence is related to heavy ion collisions experiments at RHIC and LHC [1]. The original proposal of AdS/CFT correpondence was formulated in the context of strongly coupled supersymmetric gauge theory and even more it was restricted to conformal field theory [2]. However, it is believed that at very high energy all gauge theories, at least with gravity dual, share some universalities [3]. One of the example is that the universality in the ratio between shear viscosity and entropy density [4].

A Quark Gluon Plasma (QGP) is created when two heavy ions collide at very high temperature. In this new state of matter, hadrons of the two heavy ions are disolved into a soup of quarks and gluons. A relatively heavy particle, e.g. a heavy quark, moves randomly in QGP as it collides with the constituent of QGP, quarks and gluons. Its motion can be described by a well-known Brownian motion if its time evolution of the displacement square, $s(t) \equiv x(t) - x(0)$, is given as follows [5]

$$\langle s(t)^2 \rangle \approx \begin{cases} \frac{T}{m} t^2 & (t \ll t_{\rm relax}) \\ 2Dt & (t \gg t_{\rm relax}) \end{cases}, \tag{1.1}$$

where

$$t_{\rm relax} \sim \frac{1}{\gamma_0}, \quad \gamma_0 \equiv \int_0^\infty dt\ \gamma(t), \tag{1.2}$$

is the relaxation time of the Brownian particle, $\gamma(t)$ is the memory kernel function [6],

$$D = \frac{T}{\gamma_0 m} \tag{1.3}$$

is the diffusion constant, T is the temperature, and m is the mass of Brownian particle. At early time, $t \ll t_{\rm relax}$, the Brownian particle is in balistic regime where the particle moves with constant velocity. At the late time, $t \gg t_{\rm relax}$, the particle is in diffusive regime and undergoes a random walk after collides with many particles of plasma's constituent.

In particular the QGP that we would like to consider is strongly coupled and hence we will use AdS/CFT correspondence to compute the displacement square. In general, the AdS/CFT correspondence is a correspondence between strongly coupled gauge theory in d dimension and weakly coupled gravity theory in $d+1$ dimension, with the background metric is Asymptotically AdS. An observable, $\mathcal{O}$, of gauge theory, which is hard to be computed in strongly coupled system, can be easily computed in the corresponding weakly coupled gravity theory by solving equation of motion of the dual field, ϕ, in the gravity theory where its boundary value, $\phi(r \to \infty)$, acts as a source for the observable of gauge theory [7, 8],

$$\langle e^{\int_{\partial \mathcal{M}} \mathcal{O} \phi_0} \rangle = e^{\int_{\mathcal{M}} \mathcal{L}^{\rm gravity}_{\rm on-shell}[\phi(r \to \infty) \equiv \phi_0]}, \tag{1.4}$$

with the boundary $\partial \mathcal{M}$ where the gauge theory lives in is at $r \to \infty$. At the present, we still don't have the corresponding gravity theory of QCD, the microscopic theory of QGP. However, with universalities of these gauge theories as mentioned above, we might be able to get some features of QGP qualitatively by computing using available gravity theories.

There are many calculations have been done with gravity theories where the corresponding plasma is not rotating. However, it is also natural to think that QGP should also posses some angular momentum, see [9] and reference there in. In this paper we would like to study the Brownian motion of an external quark in two dimensional rotating plasma where the corresponding gravity is three dimensional gravity with the background metric is BTZ black hole [10].

2. Gravity Setup

An external quark moving in a plasma can be presented by a fundamental string dangling from the boundary to the horizon of the background metric where the end of string at the boundary acts as a source for an external quark [11]. The length of this string related to the mass of the external quark. Therefore if the boundary is located near infinity, $r \to \infty$, the external quark is very heavy. The dynamic of this string is classically given by the Nambu-Goto action

$$S_{NG} = -\frac{1}{2\pi\alpha'}\int d\sigma^2\sqrt{(\dot{X}.X')^2 - \dot{X}^2X'^2}, \tag{2.1}$$

with

$$\dot{X}.X' = g_{\mu\nu}\frac{dX^\mu}{d\sigma^0}\frac{dX^\nu}{d\sigma^1}, \tag{2.2}$$

$$\dot{X}^2 = g_{\mu\nu}\frac{dX^\mu}{d\sigma^0}\frac{dX^\nu}{d\sigma^0}, \tag{2.3}$$

$$X'^2 = g_{\mu\nu}\frac{dX^\mu}{d\sigma^1}\frac{dX^\nu}{d\sigma^1}, \tag{2.4}$$

where $\sigma \equiv (\sigma^0, \sigma^1)$ is the string world sheet coordinate, X^μ is the spacetime coordinate, α' is related to the string length scale, and $g_{\mu\nu}$ is the background metric.

The background metric, $g_{\mu\nu}$, is given by BTZ black hole [10]:

$$\begin{aligned} ds^2 &= -N^2dt^2 + \frac{1}{N^2}dr^2 + r^2\left(N^\phi dt + d\phi\right)^2, \\ N^2(r) &= -M + \frac{r^2}{l^2} + \frac{J^2}{4r^2}, \qquad N^\phi(r) = -\frac{J}{2r^2}, \end{aligned} \tag{2.5}$$

with $-\infty < t < \infty$, $0 < r < \infty$ and $0 \leq \phi < 2\pi$. M and J are the conserved charges with asymptotic invariance under time displacement (mass) and rotational (angular momentum) invariance. The Hawking temperature of BTZ black hole, which is interpreted as the temperature of the plasma, is given by

$$\begin{aligned} T &= \frac{r_+^2 - r_-^2}{2\pi l^2 r_+}, \\ r_\pm &= \sqrt{\frac{Ml^2}{2} \pm \frac{1}{2}\xi}, \quad \xi = \sqrt{M^2l^4 - J^2l^2}, \end{aligned} \tag{2.6}$$

where r_- is the inner horizon and r_+ is the outer horizon. From now on we will set the length square dimension $l^2 = 1$ for convenient.

3. Small Fluctuation

The Nambu-Goto action under the BTZ black hole, with axial gauge $(\sigma^0, \sigma^1) = (t, r)$, becomes

$$S_{NG} = -\frac{1}{2\pi\alpha'}\int d\sigma^2 \left[\left(\dot{\phi} + N^{\phi}\right)^2 r^4 \phi'^2 - \left(N^{-2} + r^2\phi'^2\right)\left(-N^2 + r^2\left(\dot{\phi} + N^{\phi}\right)^2\right)\right]^{1/2}. \tag{3.1}$$

Under this axial gauge, we are left with one degree of freedom. So, the equation of motion for ϕ is the following:

$$-\frac{\partial}{\partial t}\left[\frac{r^2 N^{-2}}{\sqrt{-g}}\left(N^{\phi} + \dot{\phi}\right)\right] + \frac{\partial}{\partial r}\left[\frac{r^2\phi' N^2}{\sqrt{-g}}\right] = 0. \tag{3.2}$$

3.1 Trivial Solution

The trivial solution to the equation of motion (3.2) is given by $\phi(\sigma) = C$ where C is an arbitrary constant. If we take small fluctuation under this trivial solution such that $\phi = \phi + C$, $|\partial_t\phi| \ll 1$, and $|\partial_r\phi| \ll 1$, the Nambu-Goto action up to quadratic terms is given by

$$S_{NG} \approx -\frac{1}{4\pi\alpha'}\int d\sigma^2 \frac{1}{\sqrt{N^2\left(r^2 - M\right)}}\left(-\frac{N^2 r^2}{\left(r^2 - M\right)}\dot{\phi}^2 + r^2 N^4 \phi'^2\right), \tag{3.3}$$

where we have kept only the quadratic terms that contribute to equation of motion. The equation of motion of the above Nambu-Goto action is

$$-\frac{N^2 r^2}{\left(r^2 - M\right)\sqrt{N^2\left(r^2 - M\right)}}\ddot{\phi} + \frac{\partial}{\partial r}\left(r^2 N^2 \sqrt{\frac{N^2}{r^2 - M}}\phi'\right) = 0. \tag{3.4}$$

Writing the solution as $\phi(t, r) = e^{-i\omega t} f_{\omega}(r)$ and changing coordinates to $s = r^2 - \frac{1}{2}M$, the equation of motion now becomes

$$\partial_s^2 f_{\omega} + \left(\frac{3}{2(s-Q)} + \frac{3}{2(s+Q)} - \frac{1}{2(s-P)}\right)\partial_s f_{\omega} + \frac{\omega^2}{4(s-Q)(s+Q)(s-P)} f_{\omega} = 0, \tag{3.5}$$

with $P = \frac{1}{2}M$ and $Q^2 = \frac{1}{4}\xi^2$. In general, the solution to this equation is very complicated. However for the extremal case, with $|J| = M$ or $Q = 0$, the solution can be found analytically and it is given by

$$f_{\omega}^{(\pm)} = s^{-1\mp D}{}_2F_1(-1 \mp D, \frac{1}{2} \mp D; 1 \mp 2D; \frac{s}{P}), \tag{3.6}$$

with $D = \frac{1}{P}\sqrt{P\left(P + \frac{1}{4}\omega^2\right)}$ and ${}_2F_1$ is hypergeometric function. Unfortunately, this hypergeometric solution is not oscillatory function which is supposed to be the solution of this small fluctuation field ϕ or in another words this solution oscillates wildly and so it is not stable.

3.2 Linear Solution

A non-trivial solution is given by the following linear ansatz

$$\phi(t,r) = wt + \eta(r), \tag{3.7}$$

where w is a constant angular velocity. Substitute this into equation (3.2) the solution for η is

$$\eta'(r) = -\frac{\pi_\phi}{r^2N^2}\sqrt{\frac{r^4\left(w+N^\phi\right)^2 - r^2N^2}{\pi_\phi^2 - r^2N^2}}, \tag{3.8}$$

where π_ϕ is a constant related to momentum conjugate of ϕ in r direction and is also interpreted as the total force experienced by the quark [12, 13]. The negative sign in (3.8) is chosen to describe a drag force. In order to have physical solution for η, it turns out that we must fix $\pi_\phi^2 = r_{NH}^2 N(r_{NH})^2$, which is a positive number, where $r_{NH}^2 = \frac{(M-Jw)}{1-w^2}$.

This non-trivial solution also has static solution at $w = 0$ but its not equal to the above trivial solution. One can check that this static solution is curved with

$$\eta'(r) = -\frac{r_+^2 r_-^2}{(r^2-r_+^2)(r^2-r_-^2)} \tag{3.9}$$

and

$$\pi_\phi^2 = r_+^2 r_-^2 = \frac{J^2}{4}. \tag{3.10}$$

Therefore the static string feels some external force caused by the rotation of the black hole as much as

$$F_{ext} = \frac{\pi_\phi}{2\pi\alpha'} = \frac{J}{4\pi\alpha'}. \tag{3.11}$$

We must extract this external force in order to get the friction coefficient for non-zero w which is given by

$$\gamma_0 m_0 = \frac{r_+^2 + r_-^2 - 2r_+r_-w}{2\pi\alpha'(1-w^2)} = \frac{r_{NH}^2}{2\pi\alpha'}. \tag{3.12}$$

In deriving the friction coefficient, we have assumed that the momentum is non-relativistic, $P_\phi = m_0 w$, where m_0 is non-relativistic mass of the external quark.

The equation of motion from the Nambu-Goto action after substituting the small fluctuation under this background, $\phi \to wt + \eta(r) + \phi$, up to quadratic terms is quite complicated. One can check that for almost all values of w the solution does not have oscillatory modes in time since the equation contains terms with first derivative in time. However, there are values of w in which the equation of motion becomes simple and they are solutions of $J - 2Mw + Jw^2 = 0$. We pick a solution $w = r_-/r_+$ since it is consistent with the constraint $w^2 < 1$. Then the equation of motion is simply given by

$$-\frac{r^2 N}{\left(N^2 - r^2(w + N^\phi)^2\right)^{3/2}}\ddot{\phi} + \frac{\partial}{\partial r}\left[\frac{r^2 N^3 \phi'}{\left(N^2 - r^2(w + N^\phi)^2\right)^{1/2}}\right] = 0. \tag{3.13}$$

Note that this value is equal to the terminal angular velocity in the drag force formula where the total force acts on the external quark is equal to zero [14]. Changing the coordinate to $s = r^2 - r_+^2$, we obtain

$$PQ\phi'' + \left(\frac{1}{2}(PQ' - QP') + PQ'\right)\phi' - \frac{1}{4}\ddot{\phi} = 0, \tag{3.14}$$

with

$$P = N^2 - r^2(w + N^\phi)^2 = \frac{4\xi r_-^2}{J^2}s, \qquad Q = r^2 N^2 = \frac{1}{}(s + \xi)s. \tag{3.15}$$

The solution is obtain by writing $\phi(t, s) = e^{-i\omega t} f_\omega(s)$ and so the equation reduce to

$$PQf_\omega'' + \left(\frac{1}{2}(PQ' - QP') + PQ'\right)f_\omega' + \frac{\omega^2}{4}f_\omega = 0. \tag{3.16}$$

Linearly independent solutions are given by

$$f_\omega^\pm(s) = s^{\pm\frac{i\chi}{\sqrt{\xi}}}{}_2F_1(\pm\frac{i\chi}{\sqrt{\xi}}, \frac{3}{2} \pm \frac{i\chi}{\sqrt{\xi}}; 1 \pm \frac{2i\chi}{\sqrt{\xi}}; -\frac{s}{\xi}), \tag{3.17}$$

with $\chi^2 = \frac{\omega^2 J^2}{16\xi r_-^2}$. Recalling that $r_+ > r_-$, one can easily check that χ is real. The explicit form of these oscillation modes are

$$f_\omega^\pm = \frac{2^{\pm\frac{2i\chi}{\sqrt{\xi}}}}{(1 \pm \frac{2i\chi}{\sqrt{\xi}})} \frac{\left(1 \pm \frac{2i\chi}{\sqrt{s+\xi}}\right)}{\left(1 + \sqrt{1 + \frac{s}{\xi}}\right)^{\pm 2\frac{i\chi}{\sqrt{\xi}}}} s^{\pm\frac{i\chi}{\sqrt{\xi}}}. \tag{3.18}$$

Asymptotic behaviour of the solutions near the outer horizon ($s \to 0$) and boundary ($s \to \infty$) is

$$f_\omega^\pm(s) \sim \begin{cases} e^{\pm i\omega s_*}, & (s \to 0) \\ \frac{(4\xi)^{\frac{\pm i\chi}{\sqrt{\xi}}}}{(1+\frac{\pm 2i\chi}{\sqrt{\xi}})}\left[1 + \frac{2\chi^2}{s} + \cdots\right], & (s \to \infty), \end{cases} \tag{3.19}$$

with $s_* = \frac{\chi}{\omega\sqrt{\xi}}\ln(s) = \frac{J}{4\xi r_-}\ln(s)$.

4. Displacement Square

In order to have a unique solution we must impose boundary conditions to our solution as in [18]. Writing the solution

$$f_\omega(s) = A\left[f_\omega^+(s) + Bf_\omega^-(s)\right], \tag{4.1}$$

with A and B are constants. After applying Neumann boundary condition near the boundary, $s = s_c$, to implement a UV-cutoff $\partial_s f_\omega|_{s=s_c}$, we obtain

$$B = \frac{(1-\frac{2i\chi}{\sqrt{\xi}})}{(1+\frac{2i\chi}{\sqrt{\xi}})}\frac{\xi + 2i\chi\sqrt{s_c+\xi}}{\xi - 2i\chi\sqrt{s_c+\xi}}\frac{(4s_c)^{\frac{2i\chi}{\sqrt{\xi}}}}{\left(1+\sqrt{\frac{s_c+\xi}{\xi}}\right)^{\frac{4i\chi}{\sqrt{\xi}}}} \equiv e^{i\theta_\omega}. \tag{4.2}$$

Hence the constant B is merely a phase. Imposing the same Neumann boundary condition near the outer horizon at $s_h = \epsilon$, $\epsilon \ll 1$, we obtain

$$B \approx \epsilon^{\frac{2i\chi}{\sqrt{\xi}}} = e^{-\frac{2i\chi}{\sqrt{\xi}}\ln(1/\epsilon)}. \tag{4.3}$$

Imposing the second boundary condition will make the value of χ discrete and it is given by $\Delta\chi = \frac{\pi\sqrt{\xi}}{\ln(1/\epsilon)}$. In terms of ω, the discreteness is

$$\Delta\omega = \frac{4\pi\xi r_-}{J}\frac{1}{\ln(1/\epsilon)}. \tag{4.4}$$

4.1 Hawking Radiation

In this section we are going to use the formula of Hawking Radiation of the string transverse modes near the outer horizon of BTZ black hole to describe the random motion of the external quark [15, 16, 17]. As explained before, we are only interested in particular value of $w = r_-/r_+$ such that the Nambu-Goto action becomes

$$S_{NG}^{(2)} = -\frac{1}{4\pi\alpha'}\int dtdr\frac{r^2N^3\phi'^2}{(N^2 - r^2(w+N^\phi)^2)^{1/2}} - \frac{r^2N\dot{\phi}^2}{(N^2 - r^2(w+N^\phi)^2)^{3/2}}. \tag{4.5}$$

Write the general solution as

$$\phi(t,r) = \sum_{\omega=-\infty}^{\infty} e^{-i\omega t}f_\omega(r), \tag{4.6}$$

$$f_\omega(r) = g_\omega\left[f_\omega^+(r) + e^{i\theta_\omega}f_\omega^-(r)\right], \tag{4.7}$$

where the discreteness of ω is given in (4.4) and g_ω is a function of ω satisfying $g_{-\omega} = g_\omega^*$.

From (3.19), near horizon ($s \sim 0$), we have

$$\phi(t, s \to 0) \sim \sum_{\omega=-\infty}^{\infty} g_\omega\left[e^{-i\omega(t-s_*)} + e^{i\theta_\omega}e^{-i\omega(t+s_*)}\right]. \tag{4.8}$$

The right hand side in the above equation is divided into two terms. The first term is the "UP" modes and the second term is the "IN" modes. Because of Hawking radiation, the "UP" modes are excited. To see how much the "UP" modes are excited, we need to look at the Nambu-Goto action near the outer horizon. The Nambu-Goto action (4.5) becomes, for $s \to 0$,

$$S_{NG}^{(2)} \sim \frac{J^2}{16 r_-^2 \pi\alpha'} \int dt ds_* \left[\left(\frac{\partial\phi}{\partial t}\right)^2 - \left(\frac{\partial\phi}{\partial s_*}\right)^2 \right]. \tag{4.9}$$

Redefining the field to $\Phi \equiv \frac{J}{2^{3/2} r_- \sqrt{\pi\alpha'}}\phi$ then the near horizon action becomes normalized free scalar field

$$S_{NG}^{(2)} \sim \frac{1}{2} \int dt ds_* \left(\dot{\Phi}^2 - \Phi'^2 \right). \tag{4.10}$$

For a canonically normalized free scalar field Φ, the mode expansions are

$$\begin{aligned} S =& \frac{1}{2} \int dt \int_0^L dx \left(\dot{\Phi}^2 - \Phi'^2 \right), \\ \Phi(t,x) =& \frac{1}{\sqrt{L}} \sum_{k=-\infty}^{\infty} \frac{1}{\sqrt{2\upsilon_k}} \left[a_k e^{i(-\upsilon_k t + kx)} + a_k^\dagger e^{i(\upsilon_k t - kx)} \right], \\ k =& \frac{2\pi n}{L}, \;\; n \in \mathbb{Z}, \;\; \upsilon_k = |k|, \;\; \left[a_k, a_l^\dagger \right] = \delta_{kl}, \\ H =& \sum_k \upsilon_k \left(a_k^\dagger a_k + \frac{1}{2} \right), \end{aligned} \tag{4.11}$$

where L is the length of the system and we are taking periodic boundary condition. Following procedure in [18], we find k is discretized as

$$\Delta k = \frac{4\pi\xi r_-}{J} \frac{1}{\ln(1/\epsilon)}. \tag{4.12}$$

Comparing this with $\Delta k = 2\pi/L$, we also obtain the effective size of the system L which is

$$L = \frac{J}{2\xi r_-} \ln(1/\epsilon). \tag{4.13}$$

Therefore, the near outer horizon solution is written as

$$\phi(t, s \to 0) = \frac{2^{3/2} r_-^{3/2}}{J^{3/2}} \sqrt{\frac{2\xi\pi\alpha'}{\ln(1/\epsilon)}} \sum_k \frac{1}{\sqrt{2\upsilon_k}} \quad \left[a_k e^{i(-\upsilon_k t + ks_*)} + a_k^\dagger e^{i(\upsilon_k t - ks_*)} \right]. \tag{4.14}$$

There are two modes in the solution: "UP" and "IN" modes. The "UP" modes are the $k > 0$ modes and the expectation value of the occupation number is given by the Bose-Einstein distribution:

$$\langle a_k^\dagger a_l \rangle = \frac{\delta_{kl}}{e^{\beta\upsilon_k} - 1}, \qquad k > 0, \tag{4.15}$$

with $\beta = \frac{2\pi r_+}{r_+^2 - r_-^2}$. The "IN" modes ($k < 0$) just fall into the black hole where the the expectation value is zero.

Comparing (4.8) and (4.14), we can write the general solution near outer horizon as k sum [18]

$$\phi(t, s \to 0) \sim \sum_{k=-\infty}^{\infty} \left[G_k e^{i(-\upsilon_k t + k s_*)} + G_k^* e^{i(\upsilon_k t - k s_*)} \right], \tag{4.16}$$

where

$$G_{k>0} = g_k, \quad G_{k<0} = g_k^* e^{-i\theta_k}, \tag{4.17}$$

with

$$g_k = \begin{cases} \frac{2^{3/2} r_-^{3/2}}{J^{3/2}} \sqrt{\frac{\xi\pi\alpha'}{\upsilon_k \ln(1/\epsilon)}} a_k & (k > 0) \\ \frac{2^{3/2} r_-^{3/2}}{J^{3/2}} \sqrt{\frac{\xi\pi\alpha'}{\upsilon_k \ln(1/\epsilon)}} e^{-i\theta_k} a_k^\dagger & (k < 0). \end{cases} \tag{4.18}$$

Because $g_k^* = g_{-k}$, this means that a_k and $a_k^\dagger = a_{-k}$ are related.

4.2 Position of the Brownian particle

The end point of ϕ coordinate at $s = R$ is given by

$$\phi_R \equiv \phi(t, s = R) = \sum_{\upsilon=-\infty}^{\infty} \frac{g_\omega e^{-i\omega t} 2^{1+\frac{2i\chi}{\sqrt{\xi}}} \left(1 - \frac{2i\chi}{\sqrt{\xi}}\right) \xi R^{\frac{i\chi}{\sqrt{\xi}}}}{\left(1 + \sqrt{1 + \frac{R}{\xi}}\right)^{\frac{2i\chi}{\sqrt{\xi}}} \left(\xi - 2i\chi\sqrt{R + \xi}\right)}. \tag{4.19}$$

Using the relation (4.18) and (4.15), we get

$$\langle g_\omega^\dagger g_{\omega'} \rangle = \delta_{\omega\omega'} \langle a_\omega^\dagger a_{\omega'} \rangle \frac{8 r_-^3 \xi \pi \alpha'}{J^3 |\omega| \ln(1/\epsilon)}. \tag{4.20}$$

Then, we can compute

$$\begin{aligned} \langle \phi_R(t) \phi_R(0) \rangle &= \sum_{\omega > 0} \frac{32 r_-^3 \xi \pi \alpha' (\xi^2 + 4\chi^2 \xi)}{J^3 \omega (\xi^2 + 4\chi^2 (R + \xi))} \frac{1}{\ln(1/\epsilon)} \left(\frac{e^{i\omega t}}{e^{\beta\omega} - 1} + \frac{e^{-i\omega t}}{1 - e^{-\beta\omega}} \right) \\ &\approx \frac{8 r_-^2 \alpha'}{J^2} \int_0^\infty \frac{d\omega}{\omega} \frac{\left(\xi^2 + \frac{J^2}{4 r_-^2} \omega^2\right)}{\left(\xi^2 + \frac{J^2}{4 r_-^2} \frac{R + \xi}{\xi} \omega^2\right)} \left(\frac{e^{i\omega t}}{e^{\beta\omega} - 1} + \frac{e^{-i\omega t}}{1 - e^{-\beta\omega}} \right). \end{aligned} \tag{4.21}$$

The integral above is divergence. We need to regularize it by normal ordering the $a, a^\dagger$ oscillator ($: a_\omega a_\omega^\dagger :\equiv: a_\omega^\dagger a_\omega :$). The normal order correlator becomes

$$\langle : \phi_R(t) \phi_R(0) : \rangle = \frac{8 r_-^2 \alpha'}{J^2} \int_0^\infty \frac{d\omega}{\omega} \frac{\left(\xi^2 + \frac{J^2}{4 r_-^2} \omega^2\right)}{\left(\xi^2 + \frac{J^2}{4 r_-^2} \frac{R + \xi}{\xi} \omega^2\right)} \left(\frac{2\cos(\omega t)}{e^{\beta\omega} - 1} \right). \tag{4.22}$$

The displacement squared, $s(t)^2 \equiv \langle[\phi_R(t) - \phi_R(0)]^2\rangle$, is

$$s_{reg}(t)^2 = \frac{16 r_-^2 \alpha'}{J^2}\left[\frac{R}{R+\xi} I_1 + \frac{\xi}{R+\xi} I_2\right], \tag{4.23}$$

with

$$I_1 = 4\int_0^\infty \frac{dx}{x(1+a^2x^2)}\frac{\sin^2(\frac{kx}{2})}{e^x - 1}, \qquad I_2 = 4\int_0^\infty \frac{dx}{x}\frac{\sin^2(\frac{kx}{2})}{e^x - 1}, \tag{4.24}$$

and we have defined

$$x = \beta\omega, \qquad k = \frac{t}{\beta}, \qquad a^2 = \frac{J^2(R+\xi)}{4r_-^2\xi^3\beta^2}. \tag{4.25}$$

The evaluation of these integrals and their behavior for $R \gg 1$ or $a \gg 1$ can be found in Appendix B of [18]. Therefore, if $R \gg 1$, $s_{reg}(t)^2$ has the following behavior:

$$s_{reg}(t)^2 = \begin{cases} \frac{16 r_-^3 \xi^{3/2}\pi\alpha'}{J^3(R+\xi)^{1/2}\beta} t^2 + O\left(\frac{1}{R+\xi}\right) & (t \ll \beta) \\ \frac{16\pi r_-^2\alpha'}{J^2\beta} t + O\left(\log\frac{t}{\beta}\right) & (t \gg \beta). \end{cases} \tag{4.26}$$

One can check that the displacement square (4.26) is consistent with the non-rotating case computed in [18] simply by setting $J = 0$ or $r_- = 0$, which also implies $w = 0$. Using this we can write[1]

$$s_{reg}(t)^2 = \begin{cases} \frac{\alpha'}{l^4\rho_c} t^2 + O\left(\frac{1}{\rho_c^2}\right) & (t \ll \beta), \\ \frac{\alpha'\beta}{\pi l^4} t + O\left(\log\frac{t}{\beta}\right) & (t \gg \beta), \end{cases} \tag{4.27}$$

with $\rho_c^2 = 1 + \frac{R}{Ml^2}$. This is similar to the one calculated in [18] by recalling that we are using ϕ coordinate which is dimensionless.

Next, we would like to write the displacement square (4.26) in terms of physical variabels. To do so, we compute the mass of external particle using the total energy and momentum of the string under the background metric (2.5) which are given by [12]

$$E = \frac{1}{2\pi\alpha'}\int dr\ \pi_t^0, \qquad p_\phi = -\frac{1}{2\pi\alpha'}\int dr\ \pi_\phi^0, \tag{4.28}$$

with

$$\pi_t^0 = \frac{(\phi')^2}{\sqrt{-g}}\left(g_{t\phi}^2 - g_{tt}g_{\phi\phi}\right) - \frac{g_{rr}}{\sqrt{-g}}\left(g_{tt} + g_{t\phi}\dot{\phi}\right), \tag{4.29}$$

$$\pi_\phi^0 = -\frac{g_{rr}}{\sqrt{-g}}\left(g_{t\phi} + g_{\phi\phi}\dot{\phi}\right). \tag{4.30}$$

---

[1] Here, we write back l^2 by examining the dimension of all variables.

For linear solution, $\phi(t,r) = wt + \eta(r)$, with $w = r_-/r_+$ and so $\eta' = 0$, we obtain

$$E = \frac{r_+}{2\pi\alpha'\xi^{1/2}}\left(\sqrt{R+\xi} - \sqrt{\xi}\right), \qquad p_\phi = \frac{r_-}{2\pi\alpha'\xi^{1/2}}\left(\sqrt{R+\xi} - \sqrt{\xi}\right). \tag{4.31}$$

The mass is then defined as

$$m_0^2 = E^2 - p_\phi^2 = \frac{1}{(2\pi\alpha')^2}\left(\sqrt{R+\xi} - \sqrt{\xi}\right)^2. \tag{4.32}$$

Writing in terms of physical mass $m_0 \approx \frac{\sqrt{R+\xi}}{2\pi\alpha'}$, for large R, and temperature T, asymptotic time behaviour of the displacement squared is

$$\hat{s}_{reg}(t)^2 \approx \begin{cases} \left(\frac{4\xi r_-^2}{J^2}\right)^{3/2} \frac{T}{m_0} t^2 & (t \ll t_c), \\ 2Dt & (t \gg t_c), \end{cases} \tag{4.33}$$

where the diffusion constant is given by

$$D = \frac{2\pi\alpha'}{r_+^2} T. \tag{4.34}$$

So, if the Sutherland-Einstein relation holds then the relaxation time of Brownian particle is given by

$$t_c = \frac{1}{\gamma_0} = \frac{2m_0\pi\alpha'}{r_+^2}. \tag{4.35}$$

One can check that the friction coefficient is the same as in (3.12) for terminal angular velocity. Hence we have shown that small fluctuation around the linear solution at the boundary of BTZ black hole, for terminal angular velocity, behaves as a Brownian particle although the result (4.33), in the balistic regime, is different by a factor of $\left(\frac{4\xi r_-^2}{J^2}\right)^{3/2}$ compared to the standard formula of (1.1).

5. Random-Random Force Correlator

The derivation of the displacement square in the previous section is rather tedious since we have to perform the integral over the whole frequency ω. This is even more complicated if we consider the charge rotating black holes or higher dimensional black holes with more than one spatial coordinates for the Brownian motion. Another way to prove the Brownian motion of a particle which is by computing low frequency limit of the correlation function of random-random force in the Langevin equation.

Asymptotic behavior of the displacement formula for Brownian motion can be derived from the Langevin equation[2]

$$\frac{dp}{dt} = -\gamma_0 p + R(t) + K(t), \tag{5.1}$$

---

[2] This standard Langevin equation can be obtained from the generalized Langevin equation by taking zero frequency limit of the memory kernel $\gamma(t)$ [18]. Although it is constant, in general, the friction coefficient can depend on momentum or velocity as shown in [12] for the boundary metric where spacetime dimension $d \neq 4$.

where γ_0 is the friction coefficient, $R(t)$ is the random force, and $K(t)$ is the external force. The random force is the fluctuation part of the Langevin equation which takes the following properties [5]:

$$\langle R(t)\rangle = 0, \qquad \langle R(t)R(t')\rangle = \rho(t-t') \tag{5.2}$$

or in frequency modes

$$\langle R(\omega)R(\omega')\rangle = 2\pi\delta(\omega-\omega')\tilde{\rho}(\omega). \tag{5.3}$$

Assuming that $\rho(t)$ is a very sharp maximum function at $t=0$ and the equipartition of energy, one will obtain [5]

$$\int_{-\infty}^{+\infty} \rho(t)dt = \lim_{\omega\to 0}\tilde{\rho}(\omega) = 2\gamma_0 mT, \tag{5.4}$$

where m is the mass of Brownian particle and T is the temperature of the fluid. A detail derivation can be read in Appendix A and we also show there that the Brownian displacement formula can be derived simply using the Langevin equation (5.1) and low frequency limit random-random force correlator (5.4). So, to proof that our system behaves as Brownian particle we just need to compute the friction coefficient γ_0, which in holographic can be extracted from the drag force formula [13, 12] or by turning on world-volume electric field [18], and the mass of Brownian particle then compute $\tilde{\rho}(0)$ and check if the relation (5.4) is satisfied.

5.1 Non-rotating BTZ black hole

As an example, we consider a non-rotating BTZ black hole with the following metric

$$ds^2 = -(r^2-M^2)dt^2 + \frac{1}{r^2-M^2}dr^2 + r^2dX^2, \tag{5.5}$$

$$T = \frac{M}{2\pi}, \tag{5.6}$$

where M the horizon and T is the Hawking temperature. Taking fluctuation over the static solution, $\dot{X} = X' = 0$, as such the collerator of two random force in frequency mode is given by [18]

$$\langle R(\omega)R(\omega')\rangle = 2\pi\delta(\omega+\omega')\left[\frac{1}{\alpha'}4\pi T^3 + \mathcal{O}(\omega)\right]. \tag{5.7}$$

Therefore we obtain that

$$\tilde{\rho}(0) = \frac{1}{\alpha'}4\pi T^3. \tag{5.8}$$

From the drag force computation in [12], we have in non-relativistic limit

$$\gamma_0 m_0 = \frac{M^2}{2\pi\alpha'.} \tag{5.9}$$

Substituting this to (5.8) and using the Hawking temperature formula, we get the same equation as in (5.4).

5.2 Rotating BTZ black hole

As we mentioned before, the fluctuation is taken over terminal angular velocity related to the zero total force configuration. This is to simplify our computation as such only the fluctuation parts is present in the Langevin equation. Explicitly one can consider the position of the external quark into two parts: classical and fluctuation part, $X = X_{class} + \delta X$, which imply to separation of the momentum into $p = p_{class} + \delta p$. Hence the Langevin equation can be written as

$$\frac{dp}{dt} = -\gamma_0 \; p_{class} + K(t) - \gamma_0 \; \delta p + R(t). \tag{5.10}$$

In the rotating case $K(t) \neq 0$ since the fluid is rotated due to the rotation of the black hole. To cancel $K(t)$, we must fixed the angular velocity such that $\gamma_0 \; p_{class} = K(t)$, or corresponds to $\pi_\phi = 0$, which is given by the terminal angular velocity $w = r_-/r_+$ and it turns out that $r_{NH}^2 \to r_+^2$ for this value of w, see section 3.2.

The action (4.5) can be derived by considering small fluctuation over a static classical solution with the following metric

$$\begin{aligned} ds^2 = &- \sqrt{N^2(N^2 - r^2(\omega + N^\phi)^2)}dt^2 + \frac{1}{\sqrt{N^2(N^2 - r^2(\omega + N^\phi)^2)}}dr^2 \\ &+ \frac{r^2N^2}{N^2 - r^2(\omega + N^\phi)^2}d\phi^2, \end{aligned} \tag{5.11}$$

or in a simple form

$$\begin{aligned} ds^2 = &- (r^2 - r_+^2)\sqrt{\frac{(r^2 - r_-^2)(r_+^2 - r_-^2)}{r^2r_+^2}}dt^2 + \frac{1}{(r^2 - r_+^2)}\sqrt{\frac{r^2r_+^2}{(r^2 - r_-^2)(r_+^2 - r_-^2)}}dr^2 \\ &+ \frac{(r^2 - r_-^2)r_+^2}{(r_+^2 - r_-^2)}d\phi^2, \end{aligned} \tag{5.12}$$

with the Hawking temperature turns out to be equal to the Hawking temperature of rotating BTZ black hole. This metric behaves differently from the non-rotating BTZ black hole at $r \to \infty$ therefore it is not AdS at the boundary. Nevertheless, lets loose a bit the requirement of AdS space at the boundary for the Holographic Brownian motion and check if we get the same result in particular for the friction coefficient derived from the drag force formula. Repeating the derivation of admittance in [19], with a careful treatment at the boundary, one will obtain

$$\gamma_0 = \lim_{\omega \to 0} \frac{1}{\mu(\omega)} = \frac{r_+^2}{2m_0\pi\alpha'} \tag{5.13}$$

which is the same as the friction coefficient (3.12), for terminal angular velocity. Therefore we can compute

$$\tilde{\rho}(0) = \frac{r_+^2T}{\pi\alpha'} = 2\gamma_0 m_0 T \tag{5.14}$$

and we once more satisfy the formula (5.4) which implies the Brownian motion of the external quark.

6. Discussion and Conclusion

The displacement square of string fluctuation at the boundary, (4.33), asymptotically showed behaviours like a Brownian particle. In comparison with the standard asymptotic behaviour, its early times gets multiplied by a factor of

$$\left(\frac{4\xi r_-^2}{J^2}\right)^{3/2}. \tag{6.1}$$

We can write this factor in terms of terminal angular velocity $w = r_-/r_+$ such that the multiplication factor becomes $(1 - w^2)^{3/2}$. Notice that the Brownian motion formula was originally derived under non-relativistic assumption, see Appendix A. Therefore in non-relativistic limit, $w \to 0$, we get back the displacement square as in (1.1)[3].

An interesting question is whether the relativistic version of the displacement square still have the same formula as in (1.1) or it gets modified by some power of Lorentz factor, $\gamma \equiv 1/\sqrt{1 - w^2}$. In relativistic, the friction coefficient computed from the drag force formula is multiplied by a factor of γ^{-1} where the momentum is defined as $P_\phi \equiv m_r w = \gamma m_0 w$. However, the quantity $\gamma_0 m_0$ does not change its form in the relativistic version since $\gamma_0 \sim m_0^{-1}$ and so the diffusion constant is given by the same formula

$$D = \frac{T}{\gamma_r m_r}, \qquad \gamma_0 m_0 = \gamma_r m_r. \tag{6.2}$$

We could then expect that the asymptotic behaviour of the displacement square in diffusive regime would be the same as in non-relativistic case (1.1). Indeed, we found that our Holographic calculation (4.33), which is basically relativistic, showed that they are the same.

However, the displacement square of our Holographic calculation in balistic regime gets multiplied by a factor of γ^{-2} after substituting $m_0 = m_r/\gamma$. Naively, we would expect that both the regimes changes in the same way. If we define an effective mass

$$m_{eff} \equiv \frac{m_0}{(1 - w^2)^{3/2}} \tag{6.3}$$

such that an effective friction coefficient defined as

$$\gamma_{eff} = \gamma_0 (1 - w^2)^{3/2} \to D = \frac{T}{\gamma_{eff} m_{eff}} \tag{6.4}$$

thus we obtain that both the regimes does not change their forms. Another reason that both regime should retain their forms is because we have computed the fluctuation in the rest frame of the string that co-rotating with the BTZ black hole. In this co-rotating frame, the string profile is a straight line that moving with a constant angular terminal velocity. So, locally the calculation is more or less similar to the case of non-rotating BTZ black hole as in [18] regardless of the effective geometry (5.12) is different from the non-rotating BTZ black hole. Hence the displacement square must have the similar form as in (1.1) with observables, such as mass and friction coefficient, are defined in their relativistic form. Therefore we are tempted to identify the effective mass as an relativistic mass. An upshot of this we found that the relativistic mass of the external quark is multiplied by the Lorentz factor γ to the power three instead of the usual γ^1, $m_r = \gamma^3 m_0$. This peculiar power of the Lorentz factor also appears in a study of the screening length in four dimensional Kerr-AdS black hole [20] though the related observable is the mass of the black holes.

---

[3] Note that since the boundary metric is two-dimensional "Minkowski", our Holographic calculation by itself is relativistic.

As a resolution to this problem, we took an approach to analyze the Brownian motion through low frequency limit of the random-random force correlator. The calculation is more simple since we only have to look at the leading order in low frequency limit of the random-random force correlator and check if it satisfies equation (5.4). Another advantage of this approach is that the formula (5.4) does not change its form in case of relativistic or, in general, under any defined effective mass, e.g. (6.3), in condition that we also redefine the friction coefficient to an effective friction coefficient, e.g. (6.4).

One could also argue that the peculiar power of Lorentz factor appears because our calculation is not fully relativistic since the Brownian motion we considered here is non-relativistic as described in Appendix A. Therefore one need to consider the relativistic Brownian motion to get the right asymptotic behaviour of the displacement square [21, 22]. Further more, we have assumed that the Brownian motion or Langevin equation is linear in our calculation while the friction coefficient (3.12) depends on angular velocity. For angular velocity dependent friction coefficient, one need to consider non-linear Brownian motion with the following non-linear Langevin equation [23]

$$m\frac{du}{dt} = -\gamma(u)u + \psi(u)\xi(t), \tag{6.5}$$

where $\gamma(u)$ is the velocity dependent friction coefficient, $\psi(u)$ is noise function, and $\xi(t)$ is the noise. There has to be some relations between friction coefficient and the noise function for a given statistic of the noise by a general fluctuation-dissipation theorem [24]. One of those relations is derived by Dubkov, Hänggi, and Goychuk in [25] such that the friction coefficient is constant only if the noise function is an additive noise source, $\psi(u) = 1$, for white Gaussian noise, $\langle\xi(t)\xi(t')\rangle \sim \delta(t - t')$, and so it will reproduce the Einstein-Sutherland relation. However, the relativistic and/or non-linear Brownian motion are beyond our current work.

Nevertheless, it is interesting to study why the power of Lorentz factor behaves unusual for observable such as mass in the compact coordinate of rotating black holes.

Another interesting thing is to check whether this behaviour still holds for higher dimensional rotating black holes and/or for charged rotating black holes. These will be explored more in our future works.

Acknowledgments

We would like to thank the organizer of CTPNP 2012,UGM-Yogyakarta, for the opportunity to present some of our work.

A. Non-relativistic Brownian Motion

A main attention in the formulation of Brownian motion has always been about computation of the mean square value of the displacement of particle. The formula was first derived by Einstein in the case of free particle for late times [26]. A generalization for all times, in our notation, was given by Ornstein and Fürth independently [27]:

$$s^2 = \frac{2kT}{\gamma_0^2 m}\left(\gamma_0 t - 1 + e^{-\gamma_0 t}\right), \tag{A.1}$$

where k is Boltzmann's constant which was set to be one in our Holographic Brownian motion. There are several ways to derived it and in this section we are going to use the momentum method following the derivation by Uhlenbeck and Ornstein [5].

Before computing the displacement square, we are first calculating the mean (square) value of velocity $\langle u\rangle$ ($\langle u^2\rangle$). Rewriting the Langevin equation (5.1) without external force as follows:

$$\frac{du}{dt} + \gamma_0 u = A(t), \tag{A.2}$$

with $A(t) = R(t)/m$, we find the solution

$$u = u_0 e^{-\gamma_0 t} + e^{-\gamma_0 t}\int_0^t e^{\gamma_0 \zeta} A(\zeta) d\zeta, \tag{A.3}$$

where u_0 is the velocity of Brownian particle at $t = 0$. Its mean value is given by

$$\langle u\rangle^{u_0} = u_0 e^{-\gamma_0 t}, \tag{A.4}$$

$$\langle u^2\rangle^{u_0} = u_0^2 e^{-2\gamma_0 t} + e^{-2\gamma_0 t}\int_0^t\int_0^t e^{\gamma_0(\zeta_0+\zeta_1)}\langle A(\zeta_0)A(\zeta_1)\rangle d\zeta_0 d\zeta_1. \tag{A.5}$$

Defining new variables $\zeta_0 + \zeta_1 = v$ and $\zeta_0 - \zeta_1 = w$, the integral in (A.5) becomes

$$\frac{1}{2}e^{-\gamma_0 t}\int_0^{2t} e^{\gamma_0 t} dv \int_{-\infty}^{+\infty} \rho(w)\ dw = \frac{\tilde{\rho}(0)}{2\gamma_0 m^2}\left(1 - e^{-2\gamma_0 t}\right). \tag{A.6}$$

Here,we have taken the second integral from $\int_{-t}^{t} dw$ to $\int_{-\infty}^{+\infty} dw$ under an assumption that $\rho(t)$ is considered as rapidly decreasing function. A requirement from equipartition theorem, at late times, the mean square of velocity must have

$$\lim_{t\to\infty} \langle u^2 \rangle^{u_0} = \frac{kT}{m} \tag{A.7}$$

and so we obtain a constant

$$\int_{-\infty}^{+\infty} \rho(w)\ dw = \tilde{\rho}(0) = 2\gamma_0 mkT. \tag{A.8}$$

Integrating (A.3) to get the position or displacement

$$s = x - x_0 = \frac{u_0}{\gamma_0}(1 - e^{-\gamma_0 t}) - \frac{e^{-\gamma_0 t}}{\gamma_0}\int_0^t e^{\gamma_0 \zeta} A(\zeta) d\zeta + \frac{1}{\gamma_0}\int_0^t A(\zeta) d\zeta, \tag{A.9}$$

where x_0 is the initial position of the Brownian particle at $t = 0$. The mean value of position is

$$\langle s \rangle^{u_0} = \frac{u_0}{\gamma_0}(1 - e^{-\gamma_0 t}), \tag{A.10}$$

$$\langle s^2 \rangle^{u_0} = \frac{\tilde{\rho}(0)}{\gamma_0^2 m^2} t + \frac{u_0^2}{\gamma_0^2}\left(1 - e^{-\gamma_0 t}\right)^2 + \frac{\tilde{\rho}(0)}{2\gamma_0^3 m^2}\left(-3 + 4e^{-\gamma_0 t} - e^{-2\gamma_0 t}\right). \tag{A.11}$$

Taking the average over initial velocity u_0, with $\overline{u_0} = 0$ and $\overline{u_0^2} = \frac{kT}{m} = \frac{\tilde{\rho}(0)}{2\gamma_0 m^2}$, the displacement becomes

$$\overline{\langle s \rangle} = 0, \tag{A.12}$$

$$\overline{\langle s^2 \rangle} = \frac{\tilde{\rho}(0)}{\gamma_0^3 m^2}\left(\gamma_0 t - 1 + e^{-\gamma_0 t}\right). \tag{A.13}$$

Hence we obtain formula for the mean square displacement at all times and so its asymptotic time behaviour is given by (1.1). In our Holographic calculation we use the natural unit $k = 1$.

References

[1] J. Casalderrey-Solana, H. Liu, D. Mateos, K. Rajagopal and U. A. Wiedemann, "Gauge/String Duality, Hot QCD and Heavy Ion Collisions," arXiv:1101.0618 [hep-th].

[2] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. **2** (1998) 231 [Int. J. Theor. Phys. **38** (1999) 1113] [arXiv:hep-th/9711200].

[3] R. A. Janik, "The Dynamics of Quark-Gluon Plasma and AdS/CFT," Lect. Notes Phys. **828**, 147 (2011) [arXiv:1003.3291 [hep-th]].

[4] P. Kovtun, D. T. Son and A. O. Starinets, “Viscosity in strongly interacting quantum field theories from black hole physics,” Phys. Rev. Lett. **94**, 111601 (2005) [hep-th/0405231].

[5] G. E. Uhlenbeck and L. S. Ornstein, “On The Theory Of The Brownian Motion,” Phys. Rev. **36**, 823 (1930).

[6] H. Mori, “Transport, collective motion, and Brownian motion,” Prog. Theor. Phys. **33**, 423 (1965).

[7] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109].

[8] E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].

[9] F. Becattini, F. Piccinini and J. Rizzo, “Angular momentum conservation in heavy ion collisions at very high energy,” Phys. Rev. C **77**, 024906 (2008) [arXiv:0711.1253 [nucl-th]].

[10] M. Banados, C. Teitelboim and J. Zanelli, “The Black hole in three-dimensional space-time,” Phys. Rev. Lett. **69**, 1849 (1992) [arXiv:hep-th/9204099].

[11] A. Karch and E. Katz, “Adding flavor to AdS / CFT,” JHEP **0206**, 043 (2002) [hep-th/0205236].

[12] C. P. Herzog, A. Karch, P. Kovtun, C. Kozcaz and L. G. Yaffe, “Energy loss of a heavy quark moving through N = 4 supersymmetric Yang-Mills plasma,” JHEP **0607**, 013 (2006) [arXiv:hep-th/0605158].

[13] S. S. Gubser, “Drag force in AdS/CFT,” Phys. Rev. D **74**, 126005 (2006) [arXiv:hep-th/0605182].

[14] A. Nata Atmaja and K. Schalm, “Anisotropic Drag Force from 4D Kerr-AdS Black Holes,” JHEP **1104**, 070 (2011) [arXiv:1012.3800 [hep-th]].

[15] A. E. Lawrence and E. J. Martinec, “Black Hole Evaporation Along Macroscopic Strings,” Phys. Rev. D **50**, 2680 (1994) [arXiv:hep-th/9312127].

[16] S. J. Rey and J. T. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity,” Eur. Phys. J. C **22**, 379 (2001) [arXiv:hep-th/9803001].

[17] R. C. Myers, A. O. Starinets and R. M. Thomson, “Holographic spectral functions and diffusion constants for fundamental matter,” JHEP **0711**, 091 (2007) [arXiv:0706.0162 [hep-th]].

[18] J. de Boer, V. E. Hubeny, M. Rangamani and M. Shigemori, “Brownian motion in AdS/CFT,” arXiv:0812.5112 [hep-th].

[19] A. N. Atmaja, J. de Boer and M. Shigemori, "Holographic Brownian Motion and Time Scales in Strongly Coupled Plasmas," arXiv:1002.2429 [hep-th].

[20] A. N. Atmaja, "Study on Screenning Length of Quark-AntiQuark Pair in a Hot Plasma of Two Dimensional Sphere," arXiv:1111.7045 [hep-th].

[21] J. Dunkel and P. Hänggi, "Relativistic Brownian motion: From a microscopic binary collision model to the Langevin equation," Phys. Rev. E **74**, 051106 (2006) [arXiv:cond-mat/0607082v4].

[22] J. Dunkel and P. Hänggi, "Relativistic Brownian motion," Phys. Rep. **471**(1), 1-73 (2009) [arXiv:0812.1996v2].

[23] B. Lindner, "The diffusion coefficient of nonlinear Brownian motion," New J. Phys. **9** 136 (2007). doi:10.1088/1367-2630/9/5/136.

[24] R. L. Stratonovich, "Nonlinear Nonequilibrium Thermodynamics I," Springer, Berlin (1992).

[25] A. A. Dubkov, P. Hänggi and I. Goychuk, "Non-linear Brownian motion: the problem of obtaining the thermal Langevin equation for a non-Gaussian bath," J. Stat. Mech. P01034 (2009). doi:10.1088/1742-5468/2009/01/P01034.

[26] A. Einstein, Ann. d. Physik **17**, 549 (1905).

[27] L. S. Ornstein. Verls. Acad. Amst. **26**, 1005 (1917); R. Fürth, Zeits. f. Physik **2**, 244 (1920).